

## **DHSS Follow-Up Review of Air Monitoring Data from the Bridgeton Landfill Area, March 9 – March 12, 2015**

The Department of Health and Senior Services (DHSS) has reviewed air quality monitoring data collected by the Department of Natural Resources (DNR) at Bridgeton Landfill from the afternoon of **March 9, 2015** to the afternoon of **March 12, 2015**.

DNR provides continuous monitoring of reduced sulfur compounds (reported as hydrogen sulfide), sulfur dioxide, carbon monoxide, and total volatile organic compounds (VOCs) at three fixed locations as well as routine, twice daily, surveillance of hydrogen sulfide, benzene, and odor levels around the entire periphery of the landfill. DHSS has reviewed both sets of data to identify potential public health concerns for short-term health effects. Generally, samples are collected near the property boundary and dispersion is expected to reduce exposure downwind of the sample locations.

### Odors

DNR reported occasional light and moderate odors at various locations during this time period. DHSS continues to recommend that during periods of objectionable odor, sensitive individuals should stay indoors as much as possible, avoid outdoor exercise, and seek medical advice for any acute symptoms. Symptoms associated with exposure to strong odors include headache, nausea, and fatigue. Symptoms generally associated with strong odors typically disappear once the odors dissipate.

### Hydrogen Sulfide and Other Reduced Sulfur Compounds

Hydrogen sulfide concentrations were below levels of public health concern during this time period. Hydrogen sulfide levels are measured by the highly sensitive Jerome meter, which detects hydrogen sulfide specifically. Reduced sulfur compounds were periodically detected by AreaRAE monitors, but previous sampling has shown that these detections are primarily due to a reduced sulfur compound with strong odor but lower toxicity.

### Sulfur Dioxide

Sulfur dioxide concentrations were below levels of public health concern during this time period.

### Benzene and Total VOCs

Benzene was not detected in ambient air at any of the surveillance locations around the landfill during this time period. There are no health-based screening values for total VOCs. However, total VOC data are used to identify the need for compound-specific sampling. To be proactive, DNR is performing weekly VOC compound-specific sampling in locations upwind and downwind of the landfill. The laboratory results are submitted for DHSS review of public health concerns and that analysis is regularly posted online.

### Carbon Monoxide

Average carbon monoxide concentrations were below levels of public health concern.

### Radiation Rates

Gamma radiation rates continue to be indistinguishable from natural background levels and were below levels of public health concern.